bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

The establishment of HP1-independent heterochromatin reveals an essential role for HP1 proteins in maintaining epigenetic memory.

Melissa Seman[1], Alexander Levashkevich[1], Ajay Larkin[1], Fengting Huang[1], Kaushik Ragunathan[1]

AFFILIATIONS

[1]Department of Biology, Brandeis University, Waltham, MA 02451 USA

ABSTRACT

H3K9 methylation (H3K9me) marks transcriptionally silent genomic regions called heterochromatin. A conserved class of HP1 proteins are critically required to establish and maintain heterochromatin. HP1 proteins bind to H3K9me, recruit factors that promote heterochromatin formation, and oligomerize to form phase-separated condensates. We do not understand how HP1 protein binding to heterochromatin establishes and maintains transcriptional silencing. Here, we demonstrate that the *S.pombe* HP1 homolog, Swi6, can be completely bypassed to establish silencing at ectopic and endogenous loci when an H3K4 methyltransferase, Set1 and an H3K14 acetyltransferase, Mst2 are deleted. Deleting Set1 and Mst2 enhances Clr4 enzymatic activity, leading to higher H3K9me levels and increased spreading. In contrast, Swi6 and its capacity to oligomerize were indispensable during epigenetic maintenance. Our results demonstrate the role of HP1 proteins in regulating histone modification crosstalk during establishment and identifies a genetically separable function in maintaining epigenetic memory.

INTRODUCTION

Chromatin organization is critical for maintaining genome stability and regulating gene expression. Nucleosomes are at the core of chromatin structure consisting of DNA wrapped around histone proteins (H2A, H2B, H3, and H4) (Luger, 1997). Histone modifications mark two major classes of chromatin states, transcriptionally active euchromatin and silent heterochromatin. H3K9 methylation (H3K9me) is associated with constitutive heterochromatin

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

regions in organisms ranging from yeast to humans (Kouzarides, 2007; Li *et al.*, 2007; Allshire and Madhani, 2018). H3K9me promotes the binding of a conserved class of chromatin-associated proteins called heterochromatin protein 1(HP1) (Bannister *et al.*, 2001; Lachner *et al.*, 2001). HP1 proteins bind to H3K9me via an N-terminal chromodomain (CD) and dimerize through interactions involving a C-terminal chromoshadow domain (CSD) (Cowieson *et al.*, 2000; Jacobs and Khorasanizadeh, 2002). HP1 proteins can serve as adapters to recruit various effector molecules to heterochromatin, including histone modification enzymes, heterochromatin initiation factors, and nucleosome remodelers (Thiru *et al.*, 2004; Fischer *et al.*, 2009). Given the multifaceted properties of HP1 proteins, we do not understand how HP1 proteins are involved in establishing and maintaining epigenetic silencing (Eissenberg and Elgin, 2014).

In addition to their recruitment functions, HP1 proteins have important structural roles in heterochromatin formation. Based on biochemical studies, HP1 proteins are thought to bind to and bridge adjacent H3K9me nucleosomes (Machida *et al.*, 2018). Additionally, HP1 proteins can oligomerize which facilitates long-range interactions across several kilobases of DNA (Canzio *et al.*, 2011, 2013; Hinde *et al.*, 2015). *In vivo*, HP1 binding is thought to drive chromatin compaction, a characteristic feature of heterochromatin, thus creating genomic structures that are refractory to RNA polymerase II activity (Schmiedeberg *et al.*, 2004; Verschure *et al.*, 2005). *In vitro* biophysical measurements support these models wherein HP1 binding to DNA forms protein-DNA complexes that exhibit a stall force of >40pN, which represents a substantial barrier to RNA polymerase II elongation (Keenen *et al.*, 2021). Although not mutually exclusive, recent studies have also shown that the oligomerization of HP1 proteins by themselves or in complex with chromatin leads to the formation of biological condensates with liquid or gel-like characteristics (Larson *et al.*, 2017; Strom *et al.*, 2017; Sanulli *et al.*, 2019). According to these models, HP1 condensates selectively include heterochromatin associated proteins while excluding proteins that promote RNA polymerase II activity. Because of these wide-ranging biophysical properties of HP1 proteins, all models of heterochromatin are premised on HP1

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

proteins being absolutely essential for transcriptional silencing (Nishibuchi and Nakayama, 2014; Larson and Narlikar, 2018; Kumar and Kono, 2020; Schoelz and Riddle, 2022).

In *Schizosaccharomyces pombe (S.pombe),* H3K9me marks constitutive heterochromatin regions, including the pericentromeric repeats flanking centromeres, telomeres that protect the ends of chromosomes, and a yeast-specific mating type locus (Grewal and Jia, 2007). The enzyme Clr4 catalyzes H3K9 methylation, which leads to the binding of two conserved HP1 proteins, Swi6 and Chp2 (Nakayama *et al.*, 2001; Sugiyama *et al.*, 2007; Motamedi *et al.*, 2008; Fischer *et al.*, 2009). Heterochromatin formation in *S.pombe* involves three distinct steps, all of which critically require Swi6 (**Figure 1A**). First, DNA or RNA sequence-dependent *de novo* **initiation** involves the site-specific recruitment of Clr4 by RNAi factors. The pericentromeric repeats rely primarily on the RNAi pathway. Swi6 is required for RNAi-dependent initiation due to its interaction with an adapter protein, Ers1, and the loss of Swi6 leads to silencing defects at the pericentromeric repeats (Rougemaille *et al.*, 2008, 2012; Hayashi *et al.*, 2012). In contrast, the mating type locus and the telomeres depend both on RNAi and DNA binding proteins (Atf1 and Pcr1 at the mating type locus; Taz1 at the telomeres) to recruit Clr4 (Jia *et al.*, 2004; Kanoh *et al.*, 2005). Second, **establishment** involves the spreading of H3K9me across a large chromosomal region of several kilobases (Partridge *et al.*, 2000; Danzer and Wallrath, 2004). Swi6 interacts with a histone deacetylase, Clr3, and is thought to promote the spreading of H3K9me and transcriptional gene silencing, presumably through its chromatin compaction and phase separation activities (Yamada *et al.*, 2005). Third, **maintenance** involves the propagation of H3K9 methylation following DNA replication (Moazed, 2011). This is an active process that involves the read-write function of the H3K9 methyltransferase, Clr4 (Audergon *et al.*, 2015; Zhang *et al.*, 2008a; Ragunathan *et al.*, 2015). Previous studies have shown that in addition to H3K9me, Swi6 can remain bound to heterochromatin at the mating type locus throughout the cell cycle, thus raising the possibility that an HP1 bound chromatin structure, in addition to H3K9me, may be essential for epigenetic

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

inheritance (Nakayama *et al.*, 2000). However, Swi6 is a highly dynamic protein that turns over from chromatin with millisecond scale kinetics (Cheutin *et al.*, 2003, 2004). How a protein that only transiently binds chromatin can promote stable epigenetic memory remains unclear.

The simple genetic outcome of deleting Swi6 disrupts every step of heterochromatin formation, encompassing initiation, establishment, and maintenance (Hall *et al.*, 2002; Volpe *et al.*, 2002). Hence, there is a need to leverage genetic tools and cellular contexts that allow us to separately interrogate the role of HP1 proteins in each step of heterochromatin formation. To uncouple the three steps of heterochromatin formation, we utilize an inducible system (Ragunathan *et al.*, 2015). In this system, a site-specific DNA binding, TetR, is fused to the enzymatic SET domain (Clr4 without its chromodomain) of the H3K9 methyltransferase, Clr4 (TetR-Clr4-I). TetR-Clr4-I binding to *10XtetO* sites placed upstream of a reporter gene (*ade6+, ura4+,* or *GFP*) leads to heterochromatin formation and reporter gene silencing. This system captures all of the salient consequences of H3K9me establishment. Additionally, after the tetracycline-mediated release of TetR-Clr4-I, we can separately investigate how heterochromatin is maintained after DNA replication. The site-specific recruitment of Clr4 enables us to bypass RNAi-dependent initiation completely. Additionally, the TetR-Clr4-I system can be placed at both an ectopic site and the endogenous *mat* locus (Wang and Moazed 2017). Hence, our inducible system uniquely allows us to test previous models of Swi6 function in both establishing and maintaining epigenetic silencing.

Surprisingly, we discovered that Swi6 could be completely bypassed at both an ectopic site and the endogenous *mat* locus during the establishment of transcriptional gene silencing in *S. pombe*. Bypassing Swi6 required deleting two major writers of histone modifications associated with transcriptionally active chromatin, the H3K4 methyltransferase, Set1, and the H3K14 acetyltransferase, Mst2 (Noma and Grewal, 2002; Reddy *et al.*, 2011). In contrast, Swi6 was essential for maintenance and mutations that disrupt Swi6 oligomerization resulted in a loss of epigenetic inheritance. Our synthetic approach enabled us to define how Swi6 shifts the

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

equilibrium of histone post-translational modifications towards a heterochromatin-like state that is necessary to maximize Clr4 enzymatic activity. Our studies reveal how HP1 proteins regulate histone modification crosstalk and have an essential structural role in epigenetic inheritance.

RESULTS

Swi6 can be bypassed in transcriptional gene silencing when either Set1 or Mst2 are deleted.

To evaluate the effects of Swi6 downstream of H3K9 methylation, we used a synthetic approach where we tether a TetR-Clr4-I fusion protein at *10XtetO* binding sites placed upstream of a reporter gene (**Figure 1B**). As previously shown, TetR-Clr4-I recruitment was sufficient to induce heterochromatin and silence an *ade6+* reporter placed downstream of the *10XtetO* binding sequences **(Figure 1C**, *swi6+*). When the *ade6+* reporter was silenced, cells turned red when plated on media containing low adenine (Allshire *et al.*, 1994). In contrast, when *ade6+* was expressed, cells remained white. As expected, deleting Swi6 in cells where we tether Clr4 led to the loss of epigenetic silencing, as indicated by white colony color on limiting adenine media (**Figure 1C**, *swi6*Δ).

Heterochromatin is characterized by a unique repertoire of histone modifications. This includes the presence of H3K9 methylation, the absence of histone acetylation, and promoter specific H3K4 methylation, highlighting the crosstalk across different histone modifications. Therefore, we wondered if altering the existing pattern of histone modifications within a transcriptionally active region could restore silencing in cells lacking Swi6. To test whether we could reverse transcriptional activation and restore epigenetic silencing, we deleted Mst2, the catalytic subunit of a MYST acetyltransferase complex associated with H3K14 acetylation, or Set1, the catalytic subunit of the Set1C H3K4 methyltransferase complex, in *swi6*Δ cells (**Figure 1C**, *swi6*Δ*mst2*Δ or *swi6*Δ*set1*Δ*)*. We observed either no rescue upon deleting Mst2 (*swi6*Δ*mst2*Δ) or a limited rescue of *ade6+* silencing upon deleting Set1 (*swi6*Δ*set1*Δ*)*. In contrast, deleting both Set1 and Mst2 had an additive effect causing colonies to turn red

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

consistent with the full restoration of *ade6+* silencing in *swi6*Δ cells (**Figure 1C**, *swi6*Δ*set1*Δ*mst2*Δ).

To determine whether our results were generalizable to other promoters and reporter systems, we investigated silencing in strains with a *ura4-GFP* reporter whose expression depends on a short 114bp *ura4+* promoter fragment (Ragunathan *et al.*, 2015). Silencing of the GFP reporter correlates with a low fluorescence state (GFP-OFF), and GFP expression correlates with a high fluorescence state (GFP-ON). As we have previously shown, TetR-Clr4-I recruitment to *10XtetO* binding sites placed upstream of a *ura4-GFP* reporter led to low GFP expression (**Figure 1D**, *swi6+)*. Deleting Swi6 caused cells in a GFP-OFF state to shift to a GFP-ON state (**Figure 1D**, *swi6*Δ). In the GFP reporter strains, deleting Set1 in the absence of Swi6 (**Figure 1D**, *swi6*Δ*set1*Δ) partially rescued silencing while deleting Mst2 resulted in a full rescue of GFP silencing (**Figure 1D**, *swi6*Δ*mst2*Δ). Combining both deletions in the absence of Swi6 (**Figure 1D**, *swi6*Δ*set1*Δ*mst2*Δ) restored silencing similarly to what we previously observed in strains with an *ade6+* reporter. Although the two reporters behave differently, likely due to differences in their promoter sequences, deleting both Set1 and Mst2 consistently produced an additive effect.

We corroborated the results of our silencing assays by measuring transcript levels using reverse transcription followed by quantitative PCR (qRT-PCR). In the *ade6+* reporter strains, we performed qRT-PCR analysis on a gene proximal to the TetR-Clr4-I binding site, *SPCC330.06c* (strains containing the *ade6+* reporter have an endogenous *ade6-M210* allele which confounded direct qRT-PCR analysis) (Shipkovenska *et al.*, 2020). Consistent with the results of our silencing assay, *swi6*Δ*set1*Δ*mst2*Δ cells fully rescued the silencing defect of *SPCC330.06c* observed in *swi6*Δ cells (**Figure 1E**). RNA levels in *swi6*Δ*mst2*Δ*set1*Δ are further reduced approximately 10-fold lower in expression relative to *swi6+.* We observed an intermediate level of *SPCC330.06c* expression in *swi6*Δ*mst2*Δ (similar gene expression in *swi6+)* and *swi6*Δ*set1*Δ (2-fold reduction in silencing relative to *swi6+*). qRT-PCR of the GFP reporter also matched the

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

trends we observed based on fluorescence intensity measurements. GFP RNA levels increased in *swi6*Δ cells, but silencing was completely rescued in *swi6*Δ*mst2*Δ and *swi6*Δ*mst2*Δ*set1*Δ relative to *swi6*+ cells (**Figure 1F**). Intermediate levels of silencing were observed in *swi6*Δ*set1*Δ (2-fold higher than *swi6*+). We measured RNA polymerase II occupancy with ChIP-qPCR of *SPCC330.06c* using an antibody that targets the large subunit of the RNA polymerase II complex (Rpb1). Rpb1 occupancy increased upon deleting Swi6 consistent with the loss of transcriptional gene silencing (Motamedi *et al.*, 2008). Rpb1 occupancy is restored to near Swi6 wild-type levels in *swi6*Δ*mst2*Δ*set1*Δ cells (**Figure 1G**).

We also confirmed that deleting other subunits of the Set1 complex subunits (Ash2, Swd1, Swd2, Swd3, and Sdc1) instead of Set1 (*swi6*Δ*mst2*Δ+*Set1 complex deletion*) or other subunits of the Mst2 complex (Nto1, Eaf6, Ptf2, and Pdp3) instead of Mst2 (*swi6*Δ*set1*Δ+*Mst2 complex deletion*) produced a similar rescue of silencing (Reddy *et al.*, 2011; Mikheyeva *et al.*, 2014). All deletions except for *swi6*Δ*mst2*Δ*swd3*Δ rescued silencing in the *ade6*+ reporter colorimetric assay (**Figure S1A-B**). We confirmed this result using qRT-PCR assays of the endogenous *SPCC330.06c* gene (**Figure S1C-D**). All Set1 and Mst2 complex deletions led to similar levels of *SPCC330.06c* silencing except for *swd3*Δ. Finally, we confirmed that deleting proteins involved in the RNAi pathway, an endoribonuclease, Dcr1 (*swi6*Δ*mst2*Δ*set1*Δ*dcr1*Δ*)*, or the small RNA binding protein, Ago1 (*swi6*Δ*mst2*Δ*set1*Δ*ago1*Δ*)*, did not affect silencing (Volpe *et al.*, 2002). Cells remained red after the deletion of Dcr1 and Ago1, suggesting that the bypass of Swi6 at the ectopic locus upon deleting Set1 and Mst2 is distinct from any role that Swi6 may have in RNAi dependent initiation (**Figure S1E**).

Following H3K9 methylation deposition, epigenetic silencing additionally depends on distinct activities of the SHREC complex, which consists of the second HP1 protein, Chp2, a chromatin remodeler, Mit1, and a histone deacetylase, Clr3 (Sugiyama *et al.*, 2007; Motamedi *et al.*, 2008; Sadaie *et al.*, 2008; Creamer *et al.*, 2014; Job *et al.*, 2016). We deleted subunits of the SHREC complex (Mit1, Clr1, Clr2, Chp2) in *swi6*Δ*mst2*Δ*set1*Δ cells and found that silencing was

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

disrupted in each case (**Figure 1H**). These observations suggest that the entire SHREC complex is required for epigenetic silencing even when we are successfully able to bypass the requirement for Swi6. We were unable to delete Clr3 in *swi6*Δ*mst2*Δ*set1*Δ, likely because of a negative genetic interaction resulting in a synthetic lethal phenotype. Finally, we asked if Chp2 can also be bypassed by deleting Set1 and Mst2. Consistent with previous work that demonstrates a separate and nonoverlapping function for Chp2 and Swi6, deleting Mst2 or Set1 or both proteins does not restore epigenetic silencing in *chp2*Δ cells (**Figure S1F**).

Swi6 independent silencing leads to increased levels of H3K9me2 and H3K9me3.

We performed differential gene expression analysis relative to *clr4*Δ, focused on genes within a ~35kb region surrounding the TetR-Clr4-I binding site. In *swi6+* cells, several genes within the 35kb window exhibit a 2-4 fold decrease in gene expression relative to *clr4*Δ cells (**Figure 2A**). Deleting Swi6 (*swi6*Δ*)* disrupted silencing, but silencing was fully restored in *swi6*Δ*mst2*Δ*set1*Δ cells across the same 35kb region. In addition, we observed a partial rescue of silencing of many genes proximal to the TetR-Clr4-I tethering site upon deleting only Set1 or Mst2 (*swi6*Δ*set1*Δ or *swi6*Δ*mst2*Δ). However, there was an additive effect upon deleting both genes (*swi6*Δ*set1*Δ*mst2*Δ). Hence, silencing can spread across a large chromosomal region even in the absence of Swi6 following TetR-Clr4-I mediated H3K9me nucleation.

We used chromatin immunoprecipitation followed by sequencing (ChIP-seq) to determine the extent of H3K9me2 and H3K9me3 spreading across the locus where TetR-Clr4 is tethered. Deleting Swi6 led to a substantial loss of H3K9me2 and H3K9me3 across the ectopic locus despite TetR-Clr4-I being bound (*swi6*Δ, **Figure 2B-C**). Hence, the ectopic locus is highly sensitive to the presence of Swi6 for establishing and spreading H3K9 methylation. When Mst2 is deleted, there was no increase in H3K9me2 and only a slight increase in H3K9me3 across the entire region (**Figure 2B-C**, *swi6*Δ*mst2*Δ). However, when Set1 only or Set1 and Mst2 were simultaneously deleted, both H3K9me2 and H3K9me3 were approximately 4-fold higher. Notably, the domain sizes and boundaries in our Swi6 rescue mutants were similar to what we

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

observed in *swi6*+ cells (**Figure S2A-B**). We validated our ChIP-seq results using ChIP-qPCR and observed a 3-fold increase in H3K9me2 and similar levels of H3K9me3 in *swi6*Δ*mst2*Δ*set1*Δ relative to *swi6*+ with the *ade6*+ reporter (**Figure 2D-E**).

Swi6 independent silencing requires sequence-specific Clr4 tethering and H3K9 methylation.

One possibility is that depleting the epigenome of active histone modifications disrupts transcription on a genome-wide scale leading to H3K9 methylation-independent silencing. To determine if the mechanism of Swi6 independent silencing in *swi6*Δ*mst2*Δ*set1*Δ strains still required H3K9 methylation, we added tetracycline to release TetR-Clr4-I from its binding site and measured subsequent changes in RNA levels (**Figure 2F**). In the absence of Clr4 tethering, we expected the complete loss of H3K9 methylation and a concomitant loss of silencing. Consistent with our hypothesis, adding tetracycline disrupted the silencing of the *ade6*+ reporter across all genotypes, causing cells that were initially red to turn white on +tet containing medium (**Figure 2G**). Additionally, qRT-PCR confirmed that RNA transcript levels of *SPCC330.06c* returned to levels comparable to that of cells with functional Mst2 and Set1 and no Clr4 tethering (**Figure 2H**).

We compared RNA-Seq data of cells grown in +tet to cells grown without tet. We observed that 33 genes were specifically downregulated in *swi6*Δ*mst2*Δ*set1*Δ when TetR-Clr4-I was bound (-tet). Of these 33 genes, 24 were within a 200kb window surrounding the TetR-Clr4-I binding site (**Figure 2I**). In addition, we observed a similar number of genes silenced in a TetR-Clr4-I dependent manner in *swi6*+*mst2*Δ*set1*Δ (**Figure S2C**). Taken together, these results suggest that Swi6 independent silencing is dependent on TetR-Clr4-I recruitment leading to H3K9 methylation rather than spurious silencing effects due to the deletion of Set1 and Mst2.

The requirement of Swi6 for transcriptional gene silencing at the mating type locus can be bypassed upon deleting Set1 and Mst2.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Swi6 plays an essential role in RNAi-mediated heterochromatin initiation, which serves as the primary mechanism for H3K9 methylation deposition at the pericentromeric repeats and the *mat* locus. RNAi-dependent initiation at the *mat* locus can be bypassed in strains where the RNAi-initiating sequence, *cenH*, is replaced with nine copies of the *tetO* binding sites (Wang and Moazed 2017). This system recapitulates key heterochromatin features at the *mat* locus, including *cis* inheritance of ON and OFF epigenetic states. Hence, TetR-Clr4-I tethering at the *mat* locus allows us to expand our investigation of the requirement for Swi6 in transcriptional gene silencing to endogenous loci in *S. pombe*.

We used phenotypic assays based on the silencing of a *ura4+* reporter placed downstream of the TetR-Clr4-I binding site. Cells can grow on media containing 5-fluoroorotic acid (FOA) only when *ura4+* is silenced. As previously shown, we observed silencing of the *ura4+* reporter in *swi6+* cells (**Figure 3A**, *swi6+*). Deleting Swi6 caused a loss of reporter gene silencing, leading to cells that failed to grow on FOA-containing media (**Figure 3A**, *swi6Δ,*). Deleting Mst2, Set1, or both proteins led to a rescue of silencing of *ura4+* reporter compared to *swi6Δ* cells as evidenced by a rescue of growth on FOA-containing media (**Figure 3A**). We quantified the results of our growth assays by measuring *ura4+* levels using qRT-PCR measurements. Indeed, deleting Set1 or Mst2 reduces *ura4+* gene expression by half. Deleting both Set1 and Mst2 had an additive effect that reduced gene expression 5-fold relative to *swi6Δ* and achieved levels comparable to wild-type *swi6+* cells (**Figure 3B**). We used ChIP-seq to map H3K9me2 and H3K9me3 patterns across the *mat* locus. We found similar levels of H3K9me2 and H3K9me3 in wild-type and *swi6Δ* cells, likely because of the RNAi-independent recruitment of Clr4 through DNA binding proteins, Pcr1 and Atf1 (**Figure 3C**) (Jia *et al.*, 2004) . In contrast, H3K9me2 and H3K9me3 spreading increased substantially across the *mat* locus relative to wild-type cells with either *swi6Δmst2Δ* or *swi6Δset1Δ*. Notably, deleting both Set1 and Mst2 also produced an additive effect on both H3K9me2 and H3K9me3 spreading across the *mat* locus.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

TetR-Clr4 tethering is required to bypass the requirement for Swi6 at RNAi-dependent heterochromatin sites.

To test whether we could bypass the requirement for Swi6 if RNAi-dependent initiation is intact *(cenH* at the mating type locus and dg/dh repeats at the pericentromeric repeats*)*, we deleted Swi6 in strains that do not have TetR-Clr4-I mediated initiation. Consistent with the essential role of Swi6 in RNAi-mediated heterochromatin formation, we were unable to rescue silencing of a *ura4+* reporter at the *mat* locus in strains that retain the centromere homology region, *cenH* or silencing of an *ade6+* reporter at the pericentromeric repeats (**Figure S3A-B**) (Allshire *et al.*, 1995; Grewal and Klar, 1996). We observed a loss of silencing across both reporter systems upon deleting Swi6 and no rescue of silencing upon deleting Set1 and Mst2 in conjunction with Swi6.

At the telomere ends, initiation of heterochromatin formation relies on the DNA binding protein Taz1 rather than RNAi (Kanoh *et al.*, 2005). The mechanism of H3K9 methylation deposition at the rDNA locus is poorly understood, but it is known to occur through an RNAi-independent pathway (Roche *et al.*, 2016). Our ChIP-seq data showed H3K9me2 and H3K9me3 levels were increased in *swi6*Δ*mst2*Δ*set1*Δ cells relative to *swi6+* or *swi6*Δ cells at TAS sequences but not *cenH*-like sequences (**Figure 3D**). Furthermore, at the rDNA locus, H3K9me2 and H3K9me3 levels were also increased in *swi6*Δ*mst2*Δ*set1*Δ relative to *swi6+* or *swi6*Δ cells (**Figure 3E**). Therefore, we concluded that the requirement for Swi6 can be bypassed at other endogenous sites of heterochromatin formation by deleting Set1 and Mst2 if heterochromatin establishment is RNAi independent or if we can subvert the RNAi dependence at specific loci via TetR-Clr4 tethering.

Deleting Set1 and Mst2 promotes Clr4 hyperactivity

Previous studies have shown that H3K4me3 inhibits Clr4 enzymatic activity while H3K14 deacetylation, followed by ubiquitination, promotes Clr4 enzymatic activity (Greenstein *et al.*, 2020; Stirpe *et al.*, 2020). This suggests the potential for crosstalk between H3K4me3 and

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

H3K14ub in regulating Clr4 enzymatic activity. Therefore, we wondered if deleting Set1 and Mst2 primes chromatin by creating an ideal chromatin substrate for Clr4 *in vivo,* thus enabling us to bypass the requirement for Swi6.

Consistent with characteristic signatures of Clr4 hyperactivity, we observed an increase of H3K9me at rDNA and telomeres in all *swi6*Δ rescue strains (**Figure 3D-E**, *swi6*Δ*mst2*Δ*, swi6*Δ*set1*Δ*, swi6*Δ*mst2*Δ*set1*Δ) (Iglesias *et al.*, 2018). We also noted a characteristic expansion of H3K9me at meiotic genes such as *mei4* and facultative H3K9me islands such as *ssm4* in *swi6*Δ*set1*Δ and *swi6*Δ*mst2*Δ*set1*Δ cells (**Figure 4A**). In addition, we also observed a *de novo* peak of H3K9me at *clr4* which has been observed under conditions of heterochromatin misregulation (**Figure 4B**)(Wang *et al.*, 2015).

We performed genome-wide analysis on all H3K9me2 and H3K9me3 peaks across *swi6+, swi6*Δ*,* and *swi6*Δ*set1*Δ*mst2*Δ cells. We called peaks found in *swi6*Δ*mst2*Δ*set1*Δ and compared their genomic coordinates to both *swi6+* and *swi6*Δ. We saw an increase of ectopic H3K9me2 and H3K9me3 peaks at several locations in the genome in *swi6*Δ*mst2*Δ*set1*Δ that were not found in *swi6+* or *swi6*Δ (**Figure 4C-D**). These results support the hypothesis that Clr4 is hyperactive when Set1 and Mst2 are deleted in the absence of Swi6.

To validate our model of enhanced Clr4 activity upon deleting Set1 and Mst2, we engineered strains with a histone H3-G13D (*hht1-G13D*) mutation by editing one of three histone H3 copies in *S.pombe*. *hht1-G13D* is a dominant negative allele that blocks Clr4 enzymatic activity, acts in a dose-sensitive manner, and disrupts heterochromatin establishment by altering H3K9 methylation density (Cutter DiPiazza *et al.*, 2021). In strains with the ectopic *10XtetO-ade6+* reporter cells that were initially red turned white upon expressing one allele of *hht1-G13D* (**Figure 4E**). However, deleting Set1 and Mst2 in a *swi6+* or *swi6*Δ background restored silencing in *hht1-G13D* expressing cells, as indicated by the appearance of red colonies across multiple independent biological replicates. We quantified RNA levels using qRT-PCR and found that *SPCC330.06c* was not sensitive to the *hht1-G13D* mutation across different

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

strain backgrounds (data not shown). Instead, we measured expression levels of a second proximal gene, *SPCC330.07c,* in which case silencing was disrupted in cells expressing one allele of *hht1-G13D* but fully restored in *swi6*Δ*mst2*Δ*set1*Δ*hht1-G13D* cells where we observed an approximately 4-fold reduction in mRNA levels. Hence, although H3 G13D attenuates Clr4 enzymatic activity and disrupts silencing, deleting Set1 and Mst2 rescues this defect. Thus, our observations suggest that the primary role of Swi6 during establishment is to recruit factors that alter histone modification crosstalk to maximize Clr4 activity.

Swi6 is essential for epigenetic inheritance of H3K9 methylation.

Given that we could successfully bypass Swi6 during heterochromatin establishment, we wanted to test whether the same genetic conditions could allow for the bypass of Swi6 in the maintenance of epigenetic silencing. Adding tetracycline triggers the dissociation of the TetR-Clr4-I initiator, which enables us to measure H3K9me maintenance separately from its sequence-dependent establishment. In strains where we established heterochromatin at an ectopic locus, epigenetic inheritance only occurs in the absence of the putative H3K9 demethylase, Epe1 (**Figure 5A**, *epe1*Δ) (Trewick *et al.*, 2007; Audergon *et al.*, 2015; Ragunathan *et al.*, 2015). Therefore, we tested whether H3K9me and epigenetic silencing could persist in *swi6*Δ*mst2*Δ*set1*Δ*epe1*Δ strains. As expected, deleting Epe1 in *swi6*Δ*mst2*Δ*set1*Δ did not affect the establishment of *ade6*+ silencing, as indicated by red colonies on media lacking tetracycline (**Figure 5A**, *epe1*Δ*swi6*Δ*mst2*Δ*set1*Δ*)*. However, unlike *swi6*+ *epe1*Δ cells that successfully maintain epigenetic silencing and exhibit a sectored phenotype when plated on +tetracycline containing medium, *swi6*Δ*mst2*Δ*set1*Δ*epe1*Δ cells turned white on tetracycline (**Figure 5A**). In addition, *swi6*Δ*mst2*Δ*set1*Δ*epe1*Δ cells exhibited a complete loss of H3K9me2 and H3K9me3 associated with the *ade6*+ reporter when grown in tetracycline containing media consistent with the loss of silencing (**Figure 5B-C**). In contrast, cells that have *swi6*+ retained substantial levels of H3K9me2 and H3K9me3.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Additionally, we used strains where TetR-Clr4-I is required for heterochromatin establishment at the *mat* locus (Wang and Moazed 2017). Plating *swi6*+ cells on +tet containing media does not disrupt silencing as shown by the robust growth of cells on FOA+tet (**Figure 5D**, *swi6*+). Hence, cells with a functional copy of Swi6 can successfully maintain epigenetic silencing without the TetR-Clr4-I initiator. In contrast, *swi6*Δ*mst2*Δ*set1*Δ cells that could successfully silence the *ura4*+ reporter when TetR-Clr4-I was bound (despite the absence of Swi6) (**Figure 5D**), failed to maintain silencing when plated on tetracycline containing media. *swi6*Δ*mst2*Δ*set1*Δ cells exhibited a near-complete loss of growth on FOA+tet. We also confirmed through ChIP-qPCR measurements that the loss of silencing on FOA+tet correlated with the loss of H3K9me2 and H3K9me3 at the *ura4*+ reporter in *swi6*Δ*mst2*Δ*set1*Δ cells although this modification was successfully maintained in the presence of *swi6*+ (**Figure 5E-F**). Hence, our results suggest that Swi6 may have a genetically separable function in maintaining epigenetic memory.

Swi6 oligomerization is essential for epigenetic inheritance of H3K9 methylation.

Previous studies have shown that HP1 proteins (including Swi6) oligomerize to form condensates that have liquid or gel-like properties (Larson *et al.*, 2017; Strom *et al.*, 2017; Sanulli *et al.*, 2019). A mutation in the chromodomain of Swi6 (Swi6-LoopX, R93A K94A) attenuates Swi6 oligomerization by disrupting CD-CD interactions, which affects phase separation properties *in vitro* (**Figure 6A**)(Canzio *et al.*, 2013). We replaced the wild-type copy of Swi6 with a Swi6-LoopX mutant in strains that have a *10XtetO-ade6*+ reporter and an Epe1 deletion (*epe1*Δ). In *swi6-LoopX* mutants, we initially observed red colonies on media lacking tetracycline and red or sectored colonies on media containing +tetracycline (**Figure 6B**). This phenotype resembled what we observed in cells that express wild-type Swi6. However, when cells were pre-grown in +tetracycline medium prior to plating, we observed that Swi6-LoopX expressing cells turned white. In contrast, cells expressing a wild-type copy of Swi6 robustly maintained silencing exhibiting red or sectored colonies. We confirmed through ChIP-qPCR

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

measurements that H3K9me2 and H3K9me3 levels are comparable in cells with a wild-type copy of Swi6 or Swi6-LoopX during establishment (-tet, **Figure 6C-D**). These results suggest that Swi6 oligomerization is not required to establish epigenetic silencing at the ectopic locus. In contrast, ChIP-qPCR measurements of Swi6 or Swi6-LoopX expressing cells during maintenance (+tet, **Figure 6C-D**) showed significant differences in H3K9me2 and H3K9me3 levels. Swi6-LoopX expressing cells exhibited a partial loss of H3K9me2 and a complete loss of H3K9me3 during maintenance. When we replaced Swi6 with Swi6-LoopX in strains with inducible heterochromatin at the *mat* locus, *swi6-LoopX* establishment and maintenance were unaffected compared to *swi6*+, as indicated by growth on FOA and FOA+tet medium (**Figure S4A**). Unlike the ectopic locus, the *mat* locus has additional DNA binding sequences and boundary elements, which may ensure that maintenance is robust in cells expressing the Swi6-LoopX mutant. It should be noted that the essential role of Swi6 in maintenance is still apparent at the *mat* locus in a *swi6*Δ or *swi6*Δ*mst2*Δ*set1*Δ background.

DISCUSSION

HP1 proteins play multifaceted roles in heterochromatin formation and epigenetic inheritance. These roles include promoting heterochromatin specific protein-protein interactions, chromatin compaction, and the spatial relocalization of heterochromatin domains to the nuclear periphery (Thiru *et al.*, 2004; Fischer *et al.*, 2009; Holla *et al.*, 2020; Keenen *et al.*, 2021). Here, we investigate the function of the major *S. pombe* HP1 protein, Swi6, and show that it has genetically separable roles in heterochromatin initiation, establishment, and maintenance (**Figure 7**). Our data suggests that Swi6 contributes to heterochromatin establishment through protein-protein interactions, rather than its inherent structural or biophysical properties. Importantly, our findings imply that HP1-mediated chromatin compaction or protein exclusion due to HP1-mediated phase separation is unlikely to be the molecular basis for heterochromatin-mediated inhibition of transcription.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

At the pericentromeric repeats and the mating type locus, Swi6 binds Ers1, which acts as an adaptor to recruit the RNA-dependent RNA polymerase complex (RDRC) involved in double-stranded RNA synthesis(Rougemaille *et al.*, 2008, 2012; Hayashi *et al.*, 2012). In this study, we bypassed the requirement for Swi6 in initiation via TetR-Clr4-I recruitment, which enabled us to investigate the consequences of HP1 function downstream of H3K9 methylation. Our results underscore the importance of using site-specific tethering and inducible systems to investigate complex, overlapping chromatin interactions in an *in vivo* context (Stanton *et al.*, 2018).

In our experiments, we found that the deletion of Set1 and Mst2 renders Swi6 dispensable for silencing at both an ectopic site and the endogenous mating type locus. HP1 proteins lack catalytic functions, so Swi6 likely recruits histone-modifying enzymes such as Lid2, an H3K4 demethylase, Clr3, an H3K14 deacetylase, and Clr6, an H3K9 deacetylase (Fischer *et al.*, 2009). Our proposed stepwise model of heterochromatin establishment involves an initial phase of Clr4 recruitment, which leads to low levels of H3K9me (**Figure 7**). Swi6 then directs the recruitment of various histone modifiers, priming chromatin to enhance Clr4 activity further.

Previous studies have shown that silencing can occur independently of Swi6 in a synthetic context where Clr4 lacking its chromodomain (Clr4ΔCD) is recruited to an ectopic site using a Gal4 fusion protein (Kagansky *et al.*, 2009). This observation suggests that the role of Swi6 is limited to recruiting RNAi factors to pericentromeric repeats, making it dispensable in the context of an RNAi-independent heterochromatin initiation system. However, other studies have shown that Swi6 has a crucial role in transcriptional gene silencing, a downstream step of H3K9 methylation deposition (Motamedi *et al.*, 2008; Sadaie *et al.*, 2008). Our system, in which we express both the TetR-Clr4DCD initiator and a full-length copy of Clr4, recapitulates the essential role of Swi6 in transcriptional gene silencing (**Figure 1C-D**). Additionally, our results support the nonoverlapping functions of the two HP1 proteins, as we were unable to bypass the role of Chp2 by deleting Set1 and Mst2 (**Figure S1F**).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

A canonical role ascribed to HP1 proteins is heterochromatin spreading wherein H3K9 methylation extends across large chromosomal domains to sites that are distal from nucleation elements (~10-100s of kilobases) (Verschure *et al.*, 2005). In the contexts where we observed Swi6 independent silencing, we could also completely restore H3K9 methylation in excess of what is present in wild-type cells. Therefore, we favor a model in which the activity of Clr4 is the primary determinant of H3K9 methylation spreading instead of requiring processive HP1-dependent mechanisms.

Recent work has highlighted that HP1 proteins form condensates that have liquid or gel-like properties *in vitro* and in cells (Larson *et al.*, 2017; Strom *et al.*, 2017; Sanulli *et al.*, 2019). Whether the material properties of HP1- chromatin complexes are most consistent with phase separation remains deeply contested since studies of mouse chromocenters have shown that HP1a foci in cells represent collapsed chromatin globules (Erdel *et al.*, 2020). Regardless, both condensate-dependent and independent models require HP1 oligomerization. Indeed, we have demonstrated that Swi6 oligomerization is not required for heterochromatin establishment through our Swi6 delete and rescue experiments (*swi6*Δ*mst2*Δ*set1*Δ) and by replacing Swi6 with a Swi6-LoopX mutant that is defective in oligomerization (**Figure 6B**) (Canzio *et al.*, 2013). Furthermore, RNA polymerase II occupancy in *swi6*Δ*mst2*Δ*set1*Δ cells is equal to that of *swi6+* cells suggesting that mechanisms other than condensate-based exclusion can also lead to the same degree of transcriptional gene silencing (**Figure 1G**).

Deleting Swi6 or expressing the Swi6 LoopX mutant selectively disrupts epigenetic inheritance (**Figure 6A**). Oligomerization could promote various functions, including bridging, compaction, or phase separation, that are important only during maintenance. For example, HP1 phase separated condensates, could selectively include chromatin factors that may have a maintenance-specific function. Conversely, oligomerization independent of phase separation could mediate long-range chromosomal interactions or promote chromatin compaction which may be essential only during maintenance. These processes may ultimately lead to an increase

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

in the density of H3K9me which is important for epigenetic inheritance (Cutter DiPiazza *et al.*, 2021). Hence, our data suggests that HP1 proteins may be essential to create H3K9me enriched chromatin networks that facilitate the re-recruitment of Clr4 to propagate stable epigenetic states.

In summary, our studies show that HP1 proteins, given their dynamic properties, primarily function as adapters that can be repurposed to recruit diverse histone modifying proteins during establishment. Our results may explain the diversification of HP1 proteins as transcriptional activators since this would simply require the evolution of new protein-protein interactions or as cross-linkers that preserve the mechanical stability of the nucleus. In addition, HP1 proteins appear to have a genetically separable structural role during maintenance. These observations are in part consistent with recent mathematical models of how proteins that affect 3-D genome structure (such as HP1 proteins as shown in our study) in combination with limiting amounts of the enzyme are essential features to build chromatin-based systems that encode epigenetic memory (Abdulla *et al.*, 2022; Owen *et al.*, 2022).

MATERIALS AND METHODS

Strains

Some strains were made as previously described using a PCR-based gene targeting approach or with CRISPR to insert point mutations or increase transformation efficiency (Bähler *et al.*, 1998; Torres-Garcia *et al.*, 2020). Other strains were made by a cross followed by random spore analysis. All strains in this study are listed in Table S1.

Chromatin immunoprecipitation (ChIP)

As previously described (Raiymbek *et al.*, 2020), 30 ml of cells were grown to late log phase (OD_{600}- 1.8-2.2) in yeast extract supplemented with adenine or yeast extract supplemented with adenine containing tetracycline (2.5µg/ml) and fixed with 1% formaldehyde for 15 min at room temperature (RT). For Rpb1 samples, cells were resuspended in PBS with 0.25% DMSO and crosslinked with 10 mM DMA for 45 minutes at RT. Then washed once with

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

PBS and resuspended in PBS to crosslink with 1% formaldehyde for 15min at RT. 130mM glycine was then added to quench the reaction and incubated for 5 min at RT. The cells were harvested by centrifugation, and washed with TBS (50 mM Tris, pH 7.6, 500 mM NaCl). Cell pellets were resuspended in 300 µl lysis buffer (50 mM HEPES-KOH, pH 7.5, 100 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, and protease inhibitors) to which 500 µL 0.5 mm glass beads were added and cell lysis was carried out by bead beating using Omni Bead Ruptor at 3000 rpm x 30sec x 10 cycles. Tubes were punctured and the flow-through was collected in a new tube by centrifugation which was subjected to sonication to obtain fragment sizes of roughly 100-500 bp long. After sonication, the extract was centrifuged for 15 min at 13000 rpm at 4°C. The soluble chromatin was then transferred to a fresh tube and normalized for protein concentration by the Bradford assay. For each normalized sample, 25µL lysate was saved as input, to which 225 µL of 1xTE/1% SDS were added (TE: 50 mM Tris pH 8.0, 1 mM EDTA). Protein A Dynabeads were preincubated with H3K9me2 antibody (Abcam, ab1220) or H3K9me3 antibody (39161, Active Motif). For the H3K9me3 antibody (C15410193, Diagenode) preincubation was done with Streptavidin beads. For each immunoprecipitation, 2 µg H3K9me2 antibody (Abcam, ab1220), 2µg H3K9me3 antibody (39161, Active Motif) or 4 µg H3K9me3 antibody (C15410193, Diagenode) coupled to 30µL beads was added to 400µL soluble chromatin (Diagenode H3K9me3 antibody was used in Figure 2E only) and the final volume of 500µL was achieved by adding lysis buffer. Samples were incubated for 3h at 4°C. The beads were collected on magnetic stands and washed 3 times with 1 ml lysis buffer and once with 1 ml TE. For eluting bound chromatin, 100µL elution buffer I (50 mM Tris pH 8.0, 10mM EDTA, 1% SDS) was added, and the samples were incubated at 65°C for 5 min. The eluate was collected and incubated with 150 µL 1xTE/0.67% SDS in the same way. Input and immunoprecipitated samples were finally incubated overnight at 65°C to reverse crosslink. 60µg glycogen, 100 µg proteinase K (Roche), 44 ul of 5M LiCl, and 250 ul of 1xTE was added to each sample, and incubation was continued at 55°C for 1h. Phenol/chloroform extraction was carried out for all the

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

samples, followed by ethanol precipitation. Immuno-precipitated DNA was resuspended in 100µL of 10 mM Tris pH 7.5 and 50 mM NaCl. ChIP experiments were analyzed using quantitative PCR with Taq polymerase and SYBR Green using a CFX Opus 384 Real-Time PCR System. All data is presented as percent enrichment relative to input.

ChIP-Seq library preparation and processing

Libraries were constructed using the manufacturer's guidelines in the NEBNext® Ultra™ II FS DNA Library Prep Kit for Illumina, using 1ng of starting material. Barcoded libraries were pooled and sequenced with next-generation sequencing. First, raw reads were demultiplexed by barcode. Then the sequences were trimmed with Trimmomatic, aligned with BWA, and normalized by reads per million (Li and Durbin, 2010; Bolger *et al.*, 2014). Then the reads were visualized with IGV. For further analysis peaks were called using MACS2 with -g 12.57e6 in broad mode with a cutoff of 0.05 (Zhang *et al.*, 2008b). Heatmaps were generated using deepTools (v3.5.1) (Ramírez *et al.*, 2016).

RNA extraction

10ml of cells were grown to late log phase (OD_{600}- 1.8-2.2) in yeast extract supplemented with adenine. Cells were resuspended in 750uL TES buffer (0.01M Tris pH7.5, 0.01M EDTA, 0.5% SDS). Immediately 750uL of acidic phenol chloroform was added and vortexed for 2 minutes. Samples were incubated at 65C for forty minutes, vortexing for 20 seconds every ten minutes. The aqueous phase was separated by centrifuging in Phase Lock tubes for 5 minutes at 13000 rpm at 4C. The aqueous phase was transferred to new tubes and ethanol precipitated. After extraction, RNA was treated with DNase. Then the RNA was cleaned up using RNeasy Mini kits (QIAGEN). cDNA was prepared using oligoDT and SuperScript III Reverse Transcriptase (Invitrogen). The cDNA was then used for qPCR with SYBR Green and Taq polymerase on a CFX OPUS 384 Real-Time PCR System. Relative RNA is quantified using ΔCT compared to *tub* levels.

RNA-seq analysis

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Libraries were prepared, and sequencing was performed commercially. Raw fastq files were aligned to the ASM294v2 reference genome using STAR and then indexed using samtools (Dobin *et al.*, 2013). Bam files were grouped by genotype replicate and differential expression analysis was performed through Defined Region Differential Seq in the USEQ program suite (Love *et al.*, 2014). After Benjamini and Hochberg multiple testing corrections, the cutoff for significant differential expression of pairwise gene comparisons was defined as a P value of <0.01 (prior to phred transformation). Volcano plots were drawn using the ggplot2 library, and heatmap was drawn using the pheatmap library, as well as the standard R library and functions.

Silencing assays

Cells were grown in 3 ml of yeast extract containing adenine (YEA) at 32°C overnight. Cells were washed twice in water and then resuspended to a concentration of 3 x 10^7 cells/ml. 5µl of ten-fold serial dilutions was spotted on indicated plates (YE, YE+tet, EMMC, EMMC+FOA, EMMC+FOA+tet). Plates were incubated for 2-5 days before photographing.

FACS analysis

Cells containing TetR-Clr4 and a *ura4-GFP* reporter were harvested in log phase. 3 x 10^7 cells/ml were washed with water and then fixed with 70% ethanol/30% sorbitol. The cells were washed three times with 1X tris-buffered saline (TBS) and resuspended in 1ml 1xTBS. GFP fluorescence was measured using a Bio-Rad ZE5 cell analyzer with wavelength 488nm. A primary gate based on physical parameters (forward and side light scatter) was set to exclude dead cells or debris. 10,000 cells were analyzed for each sample.

ACKNOWLEDGMENTS

The authors declare no competing interests. We thank Danesh Moazed for sharing fission yeast strains used in this study. We thank Nidhi Khurana and Gulzhan Raiymbek for their support in obtaining preliminary data during the initial phase of this study and Saikat Biswas for help with running samples on the flow cytometer. We thank Amanda Ames and Basila Moochickal

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Assainar for their supportive and insightful feedback regarding this study. This work was funded by NIH award R35GM137832 to KR and T32GM007315 to MS.

REFERENCES

Abdulla, AZ, Vaillant, C, and Jost, D (2022). Painters in chromatin: a unified quantitative framework to systematically characterize epigenome regulation and memory. Nucleic Acids Res 50, 9083–9104.

Allshire, RC, Javerzat, J-P, Redhead, NJ, and Cranston, G (1994). Position Effect Variegation at Fission Yeast Centromeres.

Allshire, RC, and Madhani, HD (2018). Ten principles of heterochromatin formation and function. Nat Rev Mol Cell Biol 19, 229–244.

Allshire, RC, Nimmo, ER, Ekwall, K, Javerzat, JP, and Cranston, G (1995). Mutations derepressing silent centromeric domains in fission yeast disrupt chromosome segregation. Genes Dev 9, 218–233.

Audergon, PNCB, Catania, S, Kagansky, A, Tong, P, Shukla, M, Pidoux, AL, and Allshire, RC (2015). Restricted epigenetic inheritance of H3K9 methylation. Science 348, 132–135.

Bähler, J, Wu, J-Q, Longtine, MS, Shah, NG, Mckenzie III, A, Steever, AB, Wach, A, Philippsen, P, and Pringle, JR (1998). Heterologous modules for efficient and versatile PCR-based gene targeting inSchizosaccharomyces pombe. Yeast 14, 943–951.

Bannister, AJ, Zegerman, P, Partridge, JF, Miska, EA, Thomas, JO, Allshire, RC, and Kouzarides, T (2001). Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. Nature 410, 120–124.

Bolger, AM, Lohse, M, and Usadel, B (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120.

Canzio, D et al. (2013). A conformational switch in HP1 releases auto-inhibition to drive heterochromatin assembly. Nature 496, 377–381.

Canzio, D, Chang, EY, Shankar, S, Kuchenbecker, KM, Simon, MD, Madhani, HD, Narlikar, GJ, and Al-Sady, B (2011). Chromodomain-mediated oligomerization of HP1 suggests a nucleosome-bridging mechanism for heterochromatin assembly. Mol Cell 41, 67–81.

Cheutin, T, Gorski, SA, May, KM, Singh, PB, and Misteli, T (2004). In Vivo Dynamics of Swi6 in Yeast: Evidence for a Stochastic Model of Heterochromatin. Mol Cell Biol 24, 3157–3167.

Cheutin, T, Mcnairn, AJ, Jenuwein, T, Gilbert, DM, Singh, B, and Misteli, T (2003). Maintenance of Stable Heterochromatin Domains by Dynamic HP1 Binding.

Cowieson, NP, Partridge, JF, Allshire, RC, and McLaughlin, PJ (2000). Dimerisation of a chromo shadow domain and distinctions from the chromodomain as revealed by structural analysis. Curr Biol 10, 517–525.

Creamer, KM, Job, G, Shanker, S, Neale, GA, Lin, Y-C, Bartholomew, B, and Partridge, JF (2014). The Mi-2 Homolog Mit1 Actively Positions Nucleosomes within Heterochromatin To Suppress Transcription. Mol Cell Biol.

Cutter DiPiazza, AR, Taneja, N, Dhakshnamoorthy, J, Wheeler, D, Holla, S, and Grewal, SIS (2021). Spreading and epigenetic inheritance of heterochromatin require a critical density of histone H3 lysine 9 tri-methylation. Proc Natl Acad Sci U S A 118, 1–10.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Danzer, JR, and Wallrath, LL (2004). Mechanisms of HP1-mediated gene silencing in *Drosophila*. Development 131, 3571–3580.

Dobin, A, Davis, CA, Schlesinger, F, Drenkow, J, Zaleski, C, Jha, S, Batut, P, Chaisson, M, and Gingeras, TR (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21.

Eissenberg, JC, and Elgin, SCR (2014). HP1a: a structural chromosomal protein regulating transcription. Trends Genet 30, 103–110.

Erdel, F et al. (2020). Mouse Heterochromatin Adopts Digital Compaction States without Showing Hallmarks of HP1-Driven Liquid-Liquid Phase Separation. Mol Cell.

Fischer, T, Cui, B, Dhakshnamoorthy, J, Zhou, M, Rubin, C, Zofall, M, Veenstra, TD, and Grewal, SIS (2009). Diverse roles of HP1 proteins in heterochromatin assembly and functions in fission yeast. Proc Natl Acad Sci 106, 8998–9003.

Greenstein, RA, Barrales, RR, Sanchez, NA, Bisanz, JE, Braun, S, and Al-Sady, B (2020). Set1/COMPASS repels heterochromatin invasion at euchromatic sites by disrupting Suv39/Clr4 activity and nucleosome stability. Genes Dev 34, 99–117.

Grewal, SIS, and Jia, S (2007). Heterochromatin revisited. Nat Rev Genet 8, 35–46.

Grewal, SIS, and Klar, AJS (1996). Chromosomal inheritance of epigenetic states in fission yeast during mitosis and meiosis. Cell 86, 95–101.

Hall, IM, Shankaranarayana, GD, Noma, K ichi, Ayoub, N, Cohen, A, and Grewal, SIS (2002). Establishment and maintenance of a heterochromatin domain. Science 297, 2232–2237.

Hayashi, A, Ishida, M, Kawaguchi, R, Urano, T, Murakami, Y, and Nakayama, JI (2012). Heterochromatin protein 1 homologue Swi6 acts in concert with Ers1 to regulate RNAi-directed heterochromatin assembly. Proc Natl Acad Sci U S A 109, 6159–6164.

Hinde, E, Cardarelli, F, and Gratton, E (2015). Spatiotemporal regulation of Heterochromatin Protein 1- alpha oligomerization and dynamics in live cells. Sci Rep 5, 12001.

Holla, S, Dhakshnamoorthy, J, Folco, HD, Wheeler, D, Zofall, M, Grewal Correspondence, SIS, Balachandran, V, Xiao, H, Sun, L-L, and Grewal, SIS (2020). Positioning Heterochromatin at the Nuclear Periphery Suppresses Histone Turnover to Promote Epigenetic Inheritance In Brief Article Positioning Heterochromatin at the Nuclear Periphery Suppresses Histone Turnover to Promote Epigenetic Inheritance.

Iglesias, N, Currie, MA, Jih, G, Paulo, JA, Siuti, N, Kalocsay, M, Gygi, SP, and Moazed, D (2018). Automethylation-induced conformational switch in Clr4 (Suv39h) maintains epigenetic stability. Nature 560, 504–508.

Jacobs, SA, and Khorasanizadeh, S (2002). Structure of HP1 Chromodomain Bound to a Lysine 9-Methylated Histone H3 Tail. Science 295, 2080–2083.

Jia, S, Noma, K, and Grewal, SIS (2004). RNAi-Independent Heterochromatin Nucleation by the Stress-Activated ATF/CREB Family Proteins. Science 304, 1971–1976.

Job, G, Brugger, C, Xu, T, Bañ Os Sanz, JI, Partridge, JF, and Schalch Correspondence, T (2016). SHREC Silences Heterochromatin via Distinct Remodeling and Deacetylation Modules. Mol Cell.

Kagansky, A, Folco, HD, Almeida, R, Pidoux, AL, Boukaba, A, Simmer, F, Urano, T, Hamilton, GL, and Allshire, RC (2009). Synthetic Heterochromatin Bypasses RNAi and Centromeric Repeats to Establish Functional Centromeres. Science 324, 1716–1719.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Kanoh, J, Sadaie, M, Urano, T, and Ishikawa, F (2005). Telomere Binding Protein Taz1 Establishes Swi6 Heterochromatin Independently of RNAi at Telomeres. Curr Biol 15, 1808–1819.

Keenen, MM, Brown, D, Brennan, LD, Renger, R, Khoo, H, Carlson, CR, Huang, B, Grill, SW, Narlikar, GJ, and Redding, S (2021). HP1 proteins compact DNA into mechanically and positionally stable phase separated domains. ELife 10, e64563.

Kouzarides, T (2007). Chromatin Modifications and Their Function. Cell 128, 693–705.

Kumar, A, and Kono, H (2020). Heterochromatin protein 1 (HP1): interactions with itself and chromatin components. Biophys Rev 12, 387–400.

Lachner, M, O'Carroll, D, Rea, S, Mechtler, K, and Jenuwein, T (2001). Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. Nature 410, 116–120.

Larson, AG, Elnatan, D, Keenen, MM, Trnka, MJ, Johnston, JB, Burlingame, AL, Agard, DA, Redding, S, and Narlikar, GJ (2017). Liquid droplet formation by HP1α suggests a role for phase separation in heterochromatin. Nature 547, 236–240.

Larson, AG, and Narlikar, GJ (2018). The Role of Phase Separation in Heterochromatin Formation, Function, and Regulation. Biochemistry 57, 2540–2548.

Li, B, Carey, M, and Workman, JL (2007). The Role of Chromatin during Transcription. Cell 128, 707–719.

Li, H, and Durbin, R (2010). Fast and accurate long-read alignment with Burrows–Wheeler transform. Bioinformatics 26, 589–595.

Love, MI, Huber, W, and Anders, S (2014). Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2.

Luger, K (1997). Crystal structure of the nucleosome core particle at 2.8 A˚ resolution. 389.

Machida, S, Takizawa, Y, Ishimaru, M, Sugita, Y, Sekine, S, Nakayama, J ichi, Wolf, M, and Kurumizaka, H (2018). Structural Basis of Heterochromatin Formation by Human HP1. Mol Cell 69, 385-397.e8.

Mikheyeva, IV, Grady, PJR, Tamburini, FB, Lorenz, DR, and Cam, HP (2014). Multifaceted Genome Control by Set1 Dependent and Independent of H3K4 Methylation and the Set1C/COMPASS Complex. PLoS Genet 10.

Moazed, D (2011). Mechanisms for the Inheritance of Chromatin States. Cell 146, 510–518.

Motamedi, MR, Hong, EJE, Li, X, Gerber, S, Denison, C, Gygi, S, and Moazed, D (2008). HP1 Proteins Form Distinct Complexes and Mediate Heterochromatic Gene Silencing by Nonoverlapping Mechanisms. Mol Cell 32, 778–790.

Nakayama, J, Klar, AJS, and Grewal, SIS (2000). A Chromodomain Protein, Swi6, Performs Imprinting Functions in Fission Yeast during Mitosis and Meiosis.

Nakayama, J, Rice, JC, Strahl, BD, Allis, CD, and Grewal, SIS (2001). Role of histone H3 lysine 9 methylation in epigenetic control of heterochromatin assembly. Science 292, 110–113.

Nishibuchi, G, and Nakayama, J -i. (2014). Biochemical and structural properties of heterochromatin protein 1: understanding its role in chromatin assembly. J Biochem (Tokyo) 156, 11–20.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Noma, K ichi, and Grewal, SIS (2002). Histone H3 lysine 4 methylation is mediated by Set1 and promotes maintenance of active chromatin states in fission yeast. Proc Natl Acad Sci U S A 99, 16438–16445.

Owen, JA, Osmanović, D, and Mirny, LA (2022). Design principles of 3D epigenetic memory systems, Biophysics.

Partridge, JF, Borgstrøm, B, and Allshire, RC (2000). Distinct protein interaction domains and protein spreading in a complex centromere. Genes Dev 14, 783–791.

Ragunathan, K, Jih, G, and Moazed, D (2015). Epigenetic inheritance uncoupled from sequence-specific recruitment. Science.

Raiymbek, G, An, S, Khurana, N, Gopinath, S, Larkin, A, Biswas, S, Trievel, RC, Cho, U, and Ragunathan, K (2020). An H3K9 methylation dependent protein interaction regulates the non-enzymatic function of a putative histone demethylase. ELife 9.

Ramírez, F, Ryan, DP, Grüning, B, Bhardwaj, V, Kilpert, F, Richter, AS, Heyne, S, Dündar, F, and Manke, T (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res 44, W160–W165.

Reddy, BD, Wang, Y, Niu, L, Higuchi, EC, Marguerat, SB, Bähler, J, Smith, GR, and Jia, S (2011). Elimination of a specific histone H3K14 acetyltransferase complex bypasses the RNAi pathway to regulate pericentric heterochromatin functions. Genes Dev 25, 214–219.

Roche, B, Arcangioli, B, and Martienssen, RA (2016). RNA interference is essential for cellular quiescence. Science 354, aah5651.

Rougemaille, M, Braun, S, Coyle, S, Dumesic, PA, Garcia, JF, Isaac, RS, Libri, D, Narlikar, GJ, and Madhani, HD (2012). Ers1 links HP1 to RNAi. Proc Natl Acad Sci U S A 109, 11258–11263.

Rougemaille, M, Shankar, S, Braun, S, Rowley, M, and Madhani, HD (2008). Ers1, a rapidly diverging protein essential for RNA interference-dependent heterochromatic silencing in Schizosaccharomyces pombe. J Biol Chem 283, 25770–25773.

Sadaie, M, Kawaguchi, R, Ohtani, Y, Arisaka, F, Tanaka, K, Shirahige, K, and Nakayama, J -i. (2008). Balance between Distinct HP1 Family Proteins Controls Heterochromatin Assembly in Fission Yeast. Mol Cell Biol 28, 6973–6988.

Sanulli, S, Trnka, MJ, Dharmarajan, V, Tibble, RW, Pascal, BD, Burlingame, AL, Griffin, PR, Gross, JD, and Narlikar, GJ (2019). HP1 reshapes nucleosome core to promote phase separation of heterochromatin. 390 Nat 575.

Schmiedeberg, L, Weisshart, K, Diekmann, S, Meyer zu Hoerste, G, and Hemmerich, P (2004). High- and Low-mobility Populations of HP1 in Heterochromatin of Mammalian Cells□D. Mol Biol Cell 15.

Schoelz, JM, and Riddle, NC (2022). Functions of HP1 proteins in transcriptional regulation. Epigenetics Chromatin 15, 14.

Shipkovenska, G, Durango, A, Kalocsay, M, Gygi, SP, and Moazed, D (2020). A conserved RNA degradation complex required for spreading and epigenetic inheritance of heterochromatin. ELife 9, e54341.

Stanton, BZ, Chory, EJ, and Crabtree, GR (2018). Chemically induced proximity in biology and medicine. Science 359, eaao5902.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Stirpe, A, Guidotti, N, Northall, S, Kilic, S, Hainard, A, Vadas, O, Fierz, B, and Schalch, T (2020). SUV39 SET domains mediate crosstalk of heterochromatic histone marks Title: Regulation of SET domains by H3K14ub. BioRxiv Prepr.

Strom, AR, Emelyanov, AV, Mir, M, Fyodorov, DV, Darzacq, X, and Karpen, GH (2017). Phase separation drives heterochromatin domain formation. Nature.

Sugiyama, T, Cam, HP, Sugiyama, R, Noma, K ichi, Zofall, M, Kobayashi, R, and Grewal, SIS (2007). SHREC, an Effector Complex for Heterochromatic Transcriptional Silencing. Cell 128, 491–504.

Thiru, A, Nietlispach, D, Mott, HR, Okuwaki, M, Lyon, D, Nielsen, PR, Hirshberg, M, Verreault, A, Murzina, NV, and Laue, ED (2004). Structural basis of HP1/PXVXL motif peptide interactions and HP1 localisation to heterochromatin. EMBO J 23, 489–499.

Torres-Garcia, S, Di Pompeo, L, Eivers, L, Gaborieau, B, White, SA, Pidoux, AL, Kanigowska, P, Yaseen, I, Cai, Y, and Allshire, RC (2020). SpEDIT: A fast and efficient CRISPR/Cas9 method for fission yeast. Wellcome Open Res 5, 274.

Trewick, SC, Minc, E, Antonelli, R, Urano, T, and Allshire, RC (2007). The JmjC domain protein Epe1 prevents unregulated assembly and disassembly of heterochromatin. EMBO J 26, 4670–4682.

Verschure, PJ, van der Kraan, I, de Leeuw, W, van der Vlag, J, Carpenter, AE, Belmont, AS, and van Driel, R (2005). In Vivo HP1 Targeting Causes Large-Scale Chromatin Condensation and Enhanced Histone Lysine Methylation. Mol Cell Biol 25, 4552–4564.

Volpe, TA, Kidner, C, Hall, IM, Teng, G, Grewal, SIS, and Martienssen, RA (2002). Regulation of Heterochromatic Silencing and Histone H3 Lysine-9 Methylation by RNAi. Science 297, 1833–1837.

Wang, J, Reddy, BD, and Jia, S (2015). Rapid epigenetic adaptation to uncontrolled heterochromatin spreading. ELife 4, e06179.

Wang, X, and Moazed, D (2017). DNA sequence-dependent epigenetic inheritance of gene silencing and histone H3K9 methylation. Science 356, 88–91.


Yamada, T, Fischle, W, Sugiyama, T, David Allis, C, and Grewal, SI (2005). The Nucleation and Maintenance of Heterochromatin by a Histone Deacetylase in Fission Yeast. Mol Cell 20, 173–185.

Zhang, K, Mosch, K, Fischle, W, and Grewal, SIS (2008a). Roles of the Clr4 methyltransferase complex in nucleation, spreading and maintenance of heterochromatin. Nat Struct Mol Biol 15, 381–388.

Zhang, Y et al. (2008b). Model-based Analysis of ChIP-Seq (MACS). Genome Biol 9, R137.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 1. Swi6 can be bypassed in transcriptional gene silencing when either Set1 or Mst2 are deleted.




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 1. Swi6 can be bypassed in transcriptional gene silencing when either Set1 or Mst2 are deleted.

(**A**) Schematic describing the role of Swi6 in the three steps of heterochromatin formation. **Initiation-** DNA or RNA sequence dependent *de novo* initiation involves the site-specific recruitment of Clr4 by RNAi factors. **Establishment-** spreading of H3K9me across a large chromosomal region (several kb) and transcriptional silencing. **Maintenance-** the propagation of H3K9 methylation following DNA replication. (**B**) Schematic of TetR-Clr4-I recruitment to 10XtetO binding sites placed upstream of an *ade6*+ or ura4-GFP reporter. TetR-Clr4-I recruitment leads to inducible heterochromatin establishment. (**C**) Silencing assay of *ura4*Δ*::10XtetO-ade6*+ reporter in each indicated genotype. Red cells indicate *ade6*+ silencing. Cells are plated at 10-fold serial dilutions. (**D**) Flow cytometry to examine GFP expression in *ura4*Δ*::10XtetO-ura4-GFP* reporter in each indicated genotype. Histograms shifted to the left indicate a GFP-OFF population (red) and those shifted to the right indicate a GFP-ON population (green). (**E**) qRT-PCR measurements of RNA levels of *SPCC330.06c* in each indicated genotype. Error bars indicate SD (N=3). (**F)** qRT-PCR measurements of RNA levels of *GFP*. Error bars indicate SD (N=2). (**G**) ChIP-qPCR to measure the main subunit of RNA polymerase II, Rpb1, in each indicated genotype (N=2). (**H**) Silencing assay measuring Swi6 independent silencing in the absence of subunits of the remodeling and HDAC containing SHREC complex.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 2. Silencing relies on H3K9me but not Swi6 when Set1 and Mst2 are deleted




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 2. Silencing relies on H3K9me but not Swi6 when Set1 and Mst2 are deleted. (**A**) RNA-seq data examining expression levels relative to *clr4*Δ in the indicated genotypes surrounding the TetR-Clr4-I binding site. A 35kb window surrounding the TetR-Clr4-I binding site is shown. (**B,C**) ChIP-seq of H3K9me2 (B) and H3K9me3 (C) surrounding the TetR-Clr4-I binding site in the indicated genotypes. The *10xtetO-ade6+* at the *ura4* locus is highlighted in red. A 35kb region surrounding the tethering site is shown. All samples are normalized to reads per million. Read counts are shown in reads per hundred. (**D, E**) ChIP-qPCR to measure H3K9me2 (D) or H3K9me3 (E) at the *ade6+* reporter. Error bars indicate SD (N=3). (**F**) Schematic of TetR-Clr4-I release when cells are grown in media supplemented with tetracycline. (**G**) Silencing assay of *ura4*Δ*::10XtetO-ade6+* reporter in each indicated genotype where cells are plated on +tet media (**H**) qRT-PCR measuring RNA levels at *SPCC330.06c* in the indicated genotypes after the addition of tetracycline. Error bars indicate SD (N=3). (**I**) Volcano plots comparing genes downregulated in -tet relative to +tet conditions in *swi6*Δ*mst2*Δ*set1*Δ cells. Genes proximal to the 10xtetO-ade6+ reporter are labelled in the volcano plot.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 3. Swi6 is bypassed in heterochromatin formation when two writers of active transcription, Set1 and Mst2, are deleted at an endogenous locus.




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 3. Swi6 is bypassed in heterochromatin formation when two writers of active transcription, Set1 and Mst2, are deleted at an endogenous locus. (**A**) Inducible heterochromatin establishment at the *mat* locus. *9XtetO-ura4+* gene replaces the RNAi dependent c*enH* initiating sequence at the mating type locus. Silencing of *ura4+* reporter viaTetR-Clr4-I binding in the indicated genotypes. Cells grow on FOA only when *ura4+* is silenced. (**B**) qRT-PCR to measure RNA levels of the *ura4+* reporter in the indicated genotypes. Error bars indicate SD (N=2). (**C**) ChIP-seq of H3K9me2 and H3K9me3 at the mating type locus. The *9xtetO-ura4+* locus is highlighted in red. All samples are normalized to reads per million. Read counts are shown in reads per hundred. (**D**) Schematic of the telomere consisting of TAS sequences and RNAi dependent *cenH* like sequences. ChIP-seq of H3K9me2 and H3K9me3 at the telomere. The telomere associated sequence (TAS) is highlighted in red.(**E**) ChIP-seq of H3K9me2 and H3K9m3 at rDNA L.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 4. Deleting Set1 and Mst2 promotes Clr4 hyperactivity




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 4. Deleting Set1 and Mst2 promotes Clr4 hyperactivity. (**A**) ChIP-seq of H3K9me2 and H3K9me3 at a meiotic gene *mei4* and a facultative H3K9me island *ssm4*. (**B**) ChIP-seq of H3K9me2 and H3K9me3 at *clr4*. All ChIP-seq reads are normalized to reads per million. Read counts are shown in reads per hundred. (**C, D**) Genome wide heat map of H3K9me2 peaks (C) or H3K9me3 peaks (D). Peaks were first called in *swi6*Δ*mst2*Δ*set1*Δ and compared to H3K9me peaks at the same genomic loci in *swi6*+ and *swi6*Δ. (**E**) Schematic showing the genetic outcome of replacing a single histone H3 allele with H3 G13D (*hht1-G13D*). Silencing assay of *ura4*Δ*::10XtetO-ade6*+ reporter in the indicated genotypes. (**F**) qRT-PCR to measure gene expression at *SPCC33.07c* in the indicated genotypes. Error bars indicate SD (N=2).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 5. Swi6 is essential for heterochromatin maintenance even when Set1 and Mst2 are deleted.




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 5. Swi6 is essential for heterochromatin maintenance even when Set1 and Mst2 are deleted. (**A**) Schematic of heterochromatin maintenance assay at the ectopic locus. Adding +tetracycline leads to TetR-Clr4-I dissociation enabling us to measure H3K9me maintenance independent of sequence-dependent establishment. Silencing assay of *ura4Δ::10XtetO-ade6+* reporter in each indicated genotype without or with tetracycline. (**B, C**) ChIP-qPCR of H3K9me2 (B) or H3K9me3 (C) before and after +tetracycline addition to measure maintenance associated with the *ade6+* reporter**.** Error bars indicate SD (N=2). (**D**) Schematic of heterochromatin maintenance assay at the mat locus. Adding +tetracycline leads to TetR-Clr4-I dissociation enabling us to measure H3K9me maintenance. Maintenance, in part depends on DNA sequences at the mat locus. Silencing of *ura4+* reporter via TetR-Clr4-I binding in the indicated genotypes. Plates contain +tetracycline measure heterochromatin maintenance at the *ura4+* reporter. (**E, F**) ChIP-qPCR experiments to examine H3K9me2 (E) and H3K9me3 (F) before and after +tetracycline addition to measure establishment and maintenance. Error bars indicate SD (N=2).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 6. Swi6 oligomerization is required for maintenance.




bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 6. Swi6 oligomerization is required for maintenance. (**A**) Schematic of Swi6 oligomerization. Wild type Swi6 oligomerizes through CD-CD interactions. Swi6-LoopX (Swi6 R93A K94A) disrupts Swi6 oligomerization. (**B**) Silencing assay of *ura4Δ::10XtetO-ade6+* reporter strains in the indicated genotypes. (**C,D**) ChIP-qPCR of H3K9me2 (**C**) or H3K9me3 (**D**) with +tetracycline to measure maintenance at the *ade6+* reporter. Error bars indicate SD (N=2).

Figure 7. Model: Swi6 can be bypassed during heterochromatin establishment but is indispensable during heterochromatin maintenance.

A

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.



bioRxiv preprint doi: https://doi.org/10.1101/2023.04.30.538869; this version posted April 30, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 7. Model: Swi6 can be bypassed during heterochromatin establishment but is indispensable during heterochromatin maintenance. Initiation: Swi6 is required for initiation at RNAi dependent sites of heterochromatin formation. This process can be bypassed by sequence-specific Clr4 tethering. Establishment: The initial recruitment of Clr4 to specific sites in the genome leads to low levels of H3K9me. This is followed by Swi6 binding to recruit histone modifiers that prime chromatin which in turn enhances Clr4 enzymatic activity. Swi6 priming involves the removal of H3K4me and H3K14Ac both of which are modifications catalyzed by proteins associated with active transcription- Set1 and Mst2. Swi6 can be bypassed when Set1 and Mst2 are deleted. Maintenance: Swi6 binds to and oligomerizes in complex with H3K9me chromatin. These structural features of Swi6 binding to heterochromatin promote long-range interactions that enhance H3K9me density leading to robust epigenetic inheritance. Maintenance is lost when Swi6 oligomerization is disrupted in Swi6-LoopX mutants.